

THE AUSLANDER-REITEN QUIVER OF A POINCARÉ DUALITY SPACE

PETER JØRGENSEN

ABSTRACT. In a previous paper, Auslander-Reiten triangles and quivers were introduced into algebraic topology. This paper shows that over a Poincaré duality space, each component of the Auslander-Reiten quiver is isomorphic to $\mathbb{Z}A_\infty$.

0. INTRODUCTION

In [5], the concepts of Auslander-Reiten triangles and Auslander-Reiten quivers from the representation theory of Artin algebras were introduced into algebraic topology.

The main theorem was that Auslander-Reiten triangles exist precisely over Poincaré duality spaces. On the other hand, the only concrete Auslander-Reiten quivers computed in [5] were those over spheres. An ad hoc computation showed that the Auslander-Reiten quiver over S^d , the d -dimensional sphere, consists of $d - 1$ components, each isomorphic to $\mathbb{Z}A_\infty$.

The purpose of this paper is to show that this result was no accident. Namely, let k be a field of characteristic zero, and let X be a simply connected topological space which has Poincaré duality of dimension $d \geq 2$ over k , that is, satisfies $\mathrm{Hom}_k(H^*(X; k), k) \cong H^{d-*}(X; k)$, where $H^*(X; k)$ is singular cohomology of X with coefficients in k . The main result here is the following.

Theorem 0.1. *Let C be a component of the Auslander-Reiten quiver of the category $D^c(C^*(X; k))$. Then C is isomorphic to $\mathbb{Z}A_\infty$.*

Here $C^*(X; k)$ is the singular cochain Differential Graded Algebra of X with coefficients in k , and if $D(C^*(X; k))$ is the derived category of Differential Graded left-modules over $C^*(X; k)$, then $D^c(C^*(X; k))$ denotes the full subcategory of small objects, that is, objects M for which the functor $\mathrm{Hom}(M, -)$ preserves set indexed coproducts.

The ingredients in the proof are standard. First the Riedtmann structure theorem is invoked to show that the component C has the

2000 *Mathematics Subject Classification.* 55P62, 16E45, 16G70.

Key words and phrases. Auslander-Reiten quiver, Poincaré duality, Riedtmann structure theorem, differential graded algebra, topological space.

form $\mathbb{Z}T/\Pi$ for a directed tree T and an admissible group of automorphisms $\Pi \subseteq \text{Aut}(\mathbb{Z}T)$ (section 1). Then a certain function β is constructed on C (section 3). This induces an unbounded additive function f on T forcing its underlying graph to be A_∞ , and finally an elementary argument shows that Π acts trivially (section 4). Hence C is $\mathbb{Z}A_\infty$.

In fact, I will prove a more general result than theorem 0.1. Namely, I will not restrict myself to Differential Graded Algebras (DGAs) of the form $C^*(X; k)$, but will work over an abstract DGA denoted R .

Setup 0.2. Throughout, R denotes a DGA over the field k which satisfies the following.

- (i) R is a cochain DGA, that is, $R^i = 0$ for $i < 0$.
- (ii) $R^0 = k$.
- (iii) $R^1 = 0$.
- (iv) $\dim_k R < \infty$. □

This setup is the same as in [5, sec. 3], so I can use the results about R proved in [5].

For brief introductions to Auslander-Reiten triangles, Auslander-Reiten quivers, DGAs, Differential Graded modules (DG modules), and the way they interact, I refer the reader to [5]. My notation is mostly standard and identical to the notation of [5]; however, I do want to recapitulate a few ubiquitous items.

By $D(R)$ is denoted the derived category of DG left- R -modules, and by $D^c(R)$ the full subcategory of small objects.

By R^{op} is denoted the opposite DGA of R with multiplication

$$r \stackrel{\text{op}}{\cdot} s = (-1)^{|r||s|} sr.$$

DG left- R^{op} -modules are identified with DG right- R -modules, and so $D(R^{\text{op}})$ is identified with the derived category of DG right- R -modules, and $D^c(R^{\text{op}})$ with its full subcategory of small objects.

The DG left/right- R -modules

$$k = R/R^{\geq 1} \quad \text{and} \quad DR = \text{Hom}_k(R, k)$$

are used frequently throughout.

When I wish to emphasize that I am viewing some M as either a DG left- R -module or a DG right- R -module, I do so with subscripts, writing either ${}_R M$ or M_R .

1. APPLYING THE RIEDTMANN STRUCTURE THEOREM

This section will apply the Riedtmann structure theorem to show under some conditions that a component C of the Auslander-Reiten

quiver of $D^c(R)$ has the form $\mathbb{Z}T/\Pi$ for a directed tree T and an admissible group of automorphisms $\Pi \subseteq \text{Aut}(\mathbb{Z}T)$.

Let me start by recalling some facts from [5]. The triangulated category $D^c(R)$ does not necessarily have Auslander-Reiten triangles, but if it does, then by [5, prop. 4.4(ii)] its Auslander-Reiten translation is

$$\tau(-) = \Sigma^{-1}(DR \overset{L}{\otimes}_R -). \quad (1)$$

Here Σ denotes suspension of DG modules.

When $D^c(R)$ has Auslander-Reiten triangles, τ induces a map from the Auslander-Reiten quiver of $D^c(R)$ to itself which is also referred to as the Auslander-Reiten translation and denoted τ .

The Auslander-Reiten quiver of $D^c(R)$ is then a stable translation quiver with translation τ ; see [5, sec. 2]. This is even true in the strong sense that τ is an automorphism of the Auslander-Reiten quiver, under the extra assumption that $D^c(R^{\text{op}})$ also has Auslander-Reiten triangles. This last fact was not mentioned in [5], but is proved in the following lemma.

Lemma 1.1. *Suppose that $D^c(R)$ and $D^c(R^{\text{op}})$ have Auslander-Reiten triangles.*

- (i) *The functor $\Sigma^{-1}(DR \overset{L}{\otimes}_R -)$ is an auto-equivalence of $D^c(R)$, with quasi-inverse $\Sigma(P \overset{L}{\otimes}_R -)$ for a suitable DG left/right- R -module P .*
- (ii) *The Auslander-Reiten translation τ is an automorphism of the Auslander-Reiten quiver of $D^c(R)$.*
- (iii) *For each integer $p \neq 0$, the map τ^p is without fixed points in the Auslander-Reiten quiver of $D^c(R)$.*

Proof. (i) Both $D^c(R)$ and $D^c(R^{\text{op}})$ have Auslander-Reiten triangles, so it follows from [5, cor. 5.2 and thm. 5.1 and its proof] that there are isomorphisms ${}_R(DR) \cong {}_R(\Sigma^d R)$ in $D^c(R)$ and $(DR)_R \cong (\Sigma^d R)_R$ in $D^c(R^{\text{op}})$. (Note that I do not know ${}_R(DR)_R \cong {}_R(\Sigma^d R)_R$!)

This makes it easy to check that with P equal to $\text{RHom}_R(DR, R)$, the functors $\Sigma^{-1}(DR \overset{L}{\otimes}_R -)$ and $\Sigma(P \overset{L}{\otimes}_R -)$ are quasi-inverse endofunctors on $D^c(R)$.

(ii) This clearly follows from (i).

(iii) Let M be an indecomposable object of $D^c(R)$ with vertex $[M]$ in the Auslander-Reiten quiver of $D^c(R)$. If τ^p had the fixed point $[M]$, then p applications of equation (1) would give

$$M \cong \Sigma^{-p}((DR) \overset{L}{\otimes}_R \cdots \overset{L}{\otimes}_R (DR) \overset{L}{\otimes}_R M).$$

But this is impossible, as one proves for instance by checking

$$\inf\{i \mid H^i(M) \neq 0\} \\ \neq \inf\{i \mid H^i(\Sigma^{-p}((DR) \overset{L}{\otimes}_R \cdots \overset{L}{\otimes}_R (DR) \overset{L}{\otimes}_R M)) \neq 0\},$$

for which the general formula

$$\inf\{i \mid H^i(N \overset{L}{\otimes}_R M) \neq 0\} \\ = \inf\{i \mid H^i(N) \neq 0\} + \inf\{i \mid H^i(M) \neq 0\}$$

is handy. $\square$

A loop in a quiver is an arrow that starts and ends at the same vertex.

Lemma 1.2. *Suppose that $D^c(R)$ has Auslander-Reiten triangles. In this case, the Auslander-Reiten quiver of $D^c(R)$ has no loops.*

Proof. Suppose that there were an indecomposable object M of $D^c(R)$ whose vertex $[M]$ in the Auslander-Reiten quiver of $D^c(R)$ had a loop $[M] \longrightarrow [M]$. The arrow $[M] \longrightarrow [M]$ would mean that there existed an irreducible morphism $M \xrightarrow{\mu} M$ in $D^c(R)$. Such a morphism would be non-invertible, so it would be in the radical of the local artinian ring $\text{End}_{D^c(R)}(M)$ (cf. [5, lem. 3.6]). Hence some power μ^n would be zero.

However, it is not hard to mimick the proof of [1, lem. VII.2.5] to show that if M is indecomposable and $M \xrightarrow{\mu} M$ is irreducible, and the composition $M \xrightarrow{\mu} \cdots \xrightarrow{\mu} M$ is zero, then $[M] = \tau[M]$ holds, and this is false by lemma 1.1(iii). $\square$

In the following lemma and the rest of the paper, C is a component of the Auslander-Reiten quiver viewed as a stable translation quiver, so C is closed under the Auslander-Reiten translation τ and its inverse. This implies that C is a connected stable translation quiver with translation the restriction of τ .

For the lemma's terminology of directed trees, stable translation quivers of the form $\mathbb{Z}T$, and admissible groups of automorphisms, see [2, sec. 4.15].

Lemma 1.3. *Suppose that $D^c(R)$ and $D^c(R^{\text{op}})$ have Auslander-Reiten triangles. Let C be a component of the Auslander-Reiten quiver of $D^c(R)$.*

Then there exists a directed tree T and an admissible group of automorphisms $\Pi \subseteq \text{Aut}(\mathbb{Z}T)$ so that C is isomorphic to $\mathbb{Z}T/\Pi$ as a stable translation quiver.

Proof. Lemma 1.1(ii) implies that the restriction of τ to C is an automorphism of C , and lemma 1.2 implies that C has no loops. Also, by construction, the Auslander-Reiten quiver has no multiple arrows, so neither has C .

Hence C satisfies the conditions of the Riedtmann structure theorem, [2, thm. 4.15.6], and the conclusion of the lemma follows. $\square$

2. LABELLING THE AUSLANDER-REITEN QUIVER

This section constructs a labelling of the Auslander-Reiten quiver of $D^c(R)$, that is, a pair of positive integers $(a_{m \rightarrow n}, b_{m \rightarrow n})$ for each arrow $m \rightarrow n$ in the quiver. The construction is classical, works in good cases such as when $D^c(R)$ and $D^c(R^{\text{op}})$ both have Auslander-Reiten triangles, and goes as follows.

Let M and N be indecomposable objects of $D^c(R)$ for which there is an arrow $[M] \rightarrow [N]$ in the Auslander-Reiten quiver, and let

$$M \rightarrow X \rightarrow \tau^{-1}M \rightarrow \quad \text{and} \quad \tau N \rightarrow Y \rightarrow N \rightarrow$$

be Auslander-Reiten triangles in $D^c(R)$. Here $\tau^{-1}M$ makes sense because of lemma 1.1. Note that the triangles are determined up to isomorphism by [4, prop. 3.5(i)].

The arrow $[M] \rightarrow [N]$ means that there exists an irreducible morphism $M \rightarrow N$ in $D^c(R)$, and by [4, prop. 3.5] this means that M is a direct summand of Y and that N is a direct summand of X . Let a be the multiplicity of M as a direct summand of Y and let b be the multiplicity of N as a direct summand of X . These numbers are well-defined because $D^c(R)$ is a Krull-Schmidt category by [4, par. 3.1] and [5, lem. 3.6]. Now define the labelling by equipping $[M] \rightarrow [N]$ with the pair of positive integers (a, b) .

Lemma 2.1. *Suppose that $D^c(R)$ and $D^c(R^{\text{op}})$ have Auslander-Reiten triangles. The above labelling of the Auslander-Reiten quiver of $D^c(R)$ satisfies the following.*

- (i) $a_{\tau[M] \rightarrow [N]} = b_{[N] \rightarrow [M]}$.
- (ii) $b_{\tau[M] \rightarrow [N]} = a_{[N] \rightarrow [M]}$.
- (iii) $(a_{\tau^p[M] \rightarrow \tau^p[N]}, b_{\tau^p[M] \rightarrow \tau^p[N]}) = (a_{[M] \rightarrow [N]}, b_{[M] \rightarrow [N]})$ for each integer p .

Proof. (i) Let

$$\tau N \rightarrow Y \rightarrow N \rightarrow \tag{2}$$

be an Auslander-Reiten triangle. Then $a_{\tau[M] \rightarrow [N]}$ is defined as the multiplicity of τM as a direct summand of Y .

Now, τ is given by the formula $\Sigma^{-1}(DR \overset{L}{\otimes}_R -)$ which also defines an auto-equivalence of $D^c(R)$ by lemma 1.1(i). A quasi-inverse auto-equivalence is given by $\Sigma(P \overset{L}{\otimes}_R -)$, and applying this to the Auslander-Reiten triangle (2) gives an Auslander-Reiten triangle

$$N \longrightarrow \Sigma(P \overset{L}{\otimes}_R Y) \longrightarrow \tau^{-1}N \longrightarrow .$$

Then $b_{[N] \rightarrow [M]}$ is defined as the multiplicity of M as a direct summand of $\Sigma(P \overset{L}{\otimes}_R Y)$.

Applying $\Sigma^{-1}(DR \overset{L}{\otimes}_R -)$, this multiplicity equals the multiplicity of $\Sigma^{-1}(DR \overset{L}{\otimes}_R M)$ as a direct summand of $\Sigma^{-1}(DR \overset{L}{\otimes}_R \Sigma(P \overset{L}{\otimes}_R Y))$, that is, the multiplicity of τM as a direct summand of Y . And this again equals $a_{\tau[M] \rightarrow [N]}$ by the above.

(ii) and (iii) are proved by similar means. $\square$

3. THE FUNCTION β

Let

$$\beta(M) = \dim_k \operatorname{Ext}_R(M, k) = \dim_k H(\operatorname{RHom}_R(M, k))$$

for M in $D(R)$. Observe that β is constant on each isomorphism class in $D^c(R)$, so induces a well-defined function on the vertices $[M]$ of the Auslander-Reiten quiver of $D^c(R)$. I also denote the induced function by β , so $\beta([M]) = \beta(M)$.

The purpose of this section is to study β .

Lemma 3.1. *Suppose that $D^c(R)$ and $D^c(R^{\text{op}})$ have Auslander-Reiten triangles and that the DG left- R -module ${}_R k$ is not in $D^c(R)$.*

Let N be an indecomposable object of $D^c(R)$ with vertex $[N]$ in the Auslander-Reiten quiver of $D^c(R)$.

- (i) *If $\tau N \longrightarrow Y \longrightarrow N \longrightarrow$ is an Auslander-Reiten triangle in $D^c(R)$, then $\beta(\tau N) + \beta(N) - \beta(Y) = 0$.*
- (ii) *$\beta(\tau[N]) + \beta([N]) - \sum_{[M] \rightarrow [N]} a_{[M] \rightarrow [N]} \beta([M]) = 0$, where the sum is over all arrows in the Auslander-Reiten quiver of $D^c(R)$ which end in $[N]$.*
- (iii) *$\beta(\tau[N]) = \beta([N])$.*

Proof. (i) The Auslander-Reiten triangle $\tau N \longrightarrow Y \longrightarrow N \longrightarrow$ gives a long exact sequence consisting of pieces $\operatorname{Ext}_R^i(N, k) \longrightarrow \operatorname{Ext}_R^i(Y, k) \longrightarrow \operatorname{Ext}_R^i(\tau N, k) \xrightarrow{\delta}$. Part (i) of the lemma will follow if the connecting maps δ are zero.

And they are: By [5, lem. 4.2], the Auslander-Reiten triangle is also an Auslander-Reiten triangle in $\mathbf{D}^f(R)$, the full subcategory of $\mathbf{D}(R)$ consisting of M 's with $\dim_k HM < \infty$. Moreover, no morphism $\Sigma^{-i}(\tau N) \rightarrow {}_R k$ is a section, for any such section would clearly have to be an isomorphism, but $\Sigma^{-i}(\tau N)$ is in $\mathbf{D}^c(R)$ and ${}_R k$ is not. The definition of Auslander-Reiten triangle now says that the composition $\Sigma^{-i-1}N \rightarrow \Sigma^{-i}(\tau N) \rightarrow {}_R k$ is zero, where the first arrow is a suspension of the connecting morphism from the Auslander-Reiten triangle.

In other words, the map

$$\mathrm{Hom}_{\mathbf{D}(R)}(\Sigma^{-i}(\tau N), k) \rightarrow \mathrm{Hom}_{\mathbf{D}(R)}(\Sigma^{-i-1}N, k)$$

is zero. But this map equals

$$\mathrm{Ext}_R^i(\tau N, k) \xrightarrow{\delta} \mathrm{Ext}_R^{i+1}(N, k).$$

(ii) Consider again $\tau N \rightarrow Y \rightarrow N \rightarrow$, the Auslander-Reiten triangle from (i). From [4, prop. 3.5] follows that Y is a coproduct of copies of those indecomposable objects M of $\mathbf{D}^c(R)$ for which there are irreducible morphisms $M \rightarrow N$. That is, Y is a coproduct of copies of those indecomposable objects M of $\mathbf{D}^c(R)$ for which there are arrows $[M] \rightarrow [N]$ in the Auslander-Reiten quiver of $\mathbf{D}^c(R)$. And by the definition at the beginning of section 2, the multiplicity of M as a direct summand of Y is $a_{[M] \rightarrow [N]}$. This clearly implies

$$\beta(Y) = \sum_{[M] \rightarrow [N]} a_{[M] \rightarrow [N]} \beta([M]),$$

and then (ii) follows from (i).

(iii) Since both $\mathbf{D}^c(R)$ and $\mathbf{D}^c(R^{\mathrm{op}})$ have Auslander-Reiten triangles, it follows from [5, cor. 5.2 and thm. 5.1 and its proof] that there is an isomorphism ${}_R(DR) \cong {}_R(\Sigma^d R)$ in $\mathbf{D}^c(R)$. So as k -vector spaces,

$$\mathrm{H}(\mathrm{RHom}_R(DR, k)) \cong \mathrm{H}(\mathrm{RHom}_R(\Sigma^d R, k)) \cong \Sigma^{-d}k,$$

which easily implies $\mathrm{RHom}_R(DR, k) \cong \Sigma^{-d}k$ in $\mathbf{D}(R)$. This again gives (a) in

$$\begin{aligned} \mathrm{RHom}_R(\tau N, k) &= \mathrm{RHom}_R(\Sigma^{-1}(DR \overset{\mathrm{L}}{\otimes}_R N), k) \\ &\cong \Sigma \mathrm{RHom}_R(DR \overset{\mathrm{L}}{\otimes}_R N, k) \\ &\cong \Sigma \mathrm{RHom}_R(N, \mathrm{RHom}_R(DR, k)) \\ &\stackrel{(a)}{\cong} \Sigma \mathrm{RHom}_R(N, \Sigma^{-d}k) \\ &\cong \Sigma^{1-d} \mathrm{RHom}_R(N, k), \end{aligned}$$

and taking cohomology and k -dimension shows $\beta(\tau[N]) = \beta([N])$. $\square$

The following results leave β for a while, but return to it in corollary 3.5.

There is an abelian category $\mathcal{C}(R)$ whose objects are DG left- R -modules and whose morphisms are homomorphisms of DG left- R -modules, that is, homomorphisms of graded left-modules which are compatible with the differentials. The following Harada-Sai lemma for $\mathcal{C}(R)$ is proved just like [2, lem. 4.14.1].

Lemma 3.2. *Let $F_0, \dots, F_{2^p-1}$ be indecomposable objects of $\mathcal{C}(R)$ with $\dim_k F_i \leq p$ for each i , and let*

$$F_{2^p-1} \xrightarrow{\varphi_{2^p-1}} F_{2^p-2} \xrightarrow{\varphi_{2^p-2}} \dots \xrightarrow{\varphi_1} F_0$$

be non-isomorphisms in $\mathcal{C}(R)$.

Then $\varphi_1 \circ \dots \circ \varphi_{2^p-1} = 0$.

Corollary 3.3. *Let $M_0, \dots, M_{2^p-1}$ be indecomposable objects of $\mathcal{D}^c(R)$ with $\beta(M_i) \leq \frac{p}{\dim_k R}$ for each i , and let*

$$M_{2^p-1} \xrightarrow{\mu_{2^p-1}} M_{2^p-2} \xrightarrow{\mu_{2^p-2}} \dots \xrightarrow{\mu_1} M_0$$

be non-isomorphisms in $\mathcal{D}^c(R)$.

Then $\mu_1 \circ \dots \circ \mu_{2^p-1} = 0$.

Proof. Pick minimal semi-free resolutions $F_i \xrightarrow{\simeq} M_i$ (minimality means that the differential ∂_{F_i} takes values inside $R^{\geq 1} \cdot F_i$, see [5, lem. 3.3]). Then the morphisms $M_{2^p-1} \xrightarrow{\mu_{2^p-1}} M_{2^p-2} \xrightarrow{\mu_{2^p-2}} \dots \xrightarrow{\mu_1} M_0$ in $\mathcal{D}^c(R)$ are represented by morphisms $F_{2^p-1} \xrightarrow{\varphi_{2^p-1}} F_{2^p-2} \xrightarrow{\varphi_{2^p-2}} \dots \xrightarrow{\varphi_1} F_0$ in $\mathcal{C}(R)$.

If φ_i were an isomorphism in $\mathcal{C}(R)$, then μ_i would be an isomorphism in $\mathcal{D}^c(R)$, so each φ_i is a non-isomorphism in $\mathcal{C}(R)$. Also, since F_i is minimal, it is easy to see that if F_i decomposed non-trivially in $\mathcal{C}(R)$, then M_i would decompose non-trivially in $\mathcal{D}^c(R)$, so each F_i is indecomposable in $\mathcal{C}(R)$.

Now

$$\begin{aligned} \text{Ext}_R(M_i, k) &= \text{H}(\text{RHom}_R(M_i, k)) \cong \text{H}(\text{Hom}_R(F_i, k)) \\ &\stackrel{(a)}{\cong} \text{Hom}_R(F_i, k)^\natural = \text{Hom}_{R^\natural}(F_i^\natural, k^\natural). \end{aligned} \tag{3}$$

Here $\natural$ indicates the functor which forgets differentials, and therefore sends DGAs to graded algebras and DG modules to graded modules. The isomorphism (a) holds because F_i is minimal, whence the differential of the complex $\text{Hom}_R(F_i, k)$ is zero, so taking cohomology amounts simply to forgetting the differential.

Taking k -dimensions in equation (3) shows that $\beta(M_i)$ equals the k -dimension of $\text{Hom}_{R^\natural}(F_i^\natural, k^\natural)$. However, since F_i is semi-free I have

$$F_i^\natural \cong \bigoplus_j \Sigma^{\sigma_j}(R^\natural)$$

for certain integers σ_j , and the k -dimension of $\text{Hom}_{R^\natural}(F_i^\natural, k^\natural)$ clearly equals the number of summands $\Sigma^{\sigma_j}(R^\natural)$ in $F_i^\natural$.

All in all, $\beta(M_i)$ equals the number of summands $\Sigma^{\sigma_j}(R^\natural)$ in $F_i^\natural$ which gives (b) in

$$\begin{aligned} \dim_k F_i &= \dim_k F_i^\natural \stackrel{(b)}{=} \beta(M_i) \dim_k R^\natural \\ &= \beta(M_i) \dim_k R \leq \frac{p}{\dim_k R} \dim_k R = p. \end{aligned}$$

But now lemma 3.2 gives $\varphi_1 \circ \cdots \circ \varphi_{2^p-1} = 0$ which clearly implies $\mu_i \circ \cdots \circ \mu_{2^p-1} = 0$. $\square$

Lemma 3.4. *Suppose that $D^c(R)$ has Auslander-Reiten triangles and that ${}_R k$ is not in $D^c(R)$.*

Given an indecomposable object M_0 of $D^c(R)$ and an integer $q \geq 0$, there exist indecomposable objects and irreducible morphisms

$$M_q \xrightarrow{\mu_q} M_{q-1} \xrightarrow{\mu_{q-1}} \cdots \xrightarrow{\mu_1} M_0$$

in $D^c(R)$ with $\mu_1 \circ \cdots \circ \mu_q \neq 0$.

Proof. If M is a non-zero object in $D^f(R)$, the full subcategory of $D(R)$ consisting of M 's with $\dim_k HM < \infty$, then the k -dual $DM = \text{Hom}_k(M, k)$ is non-zero in $D^f(R^{\text{op}})$, so the minimal semi-free resolution F of DM is non-trivial. Hence $\text{RHom}_{R^{\text{op}}}(DM, k) \cong \text{Hom}_{R^{\text{op}}}(F, k)$ has non-zero cohomology, so by dualization the same holds for

$$\text{RHom}_R(Dk, DDM) \cong \text{RHom}_R(k, M).$$

So there exists an i so that $H^{-i}(\text{RHom}_R(k, M)) \cong \text{Hom}_{D(R)}(\Sigma^i k, M)$ is non-zero; that is,

$$\begin{aligned} &\text{For } M \text{ non-zero in } D^f(R), \text{ there exists} \\ &\text{a non-zero morphism } {}_R(\Sigma^i k) \longrightarrow M \text{ in } D^f(R). \end{aligned} \quad (4)$$

Moreover, since M is non-zero, each retraction ${}_R(\Sigma^i k) \longrightarrow M$ in $D^f(R)$ is clearly an isomorphism. If M is in $D^c(R)$, then there is no such isomorphism because ${}_R(\Sigma^i k)$ is not in $D^c(R)$, so

$$\begin{aligned} &\text{For } M \text{ non-zero in } D^c(R), \text{ no} \\ &\text{morphism } {}_R(\Sigma^i k) \longrightarrow M \text{ is a retraction in } D^f(R). \end{aligned} \quad (5)$$

Now for the proof proper. In fact, I shall prove slightly more than claimed, namely, there exists

$${}_R(\Sigma^i k) \xrightarrow{\kappa_q} M_q \xrightarrow{\mu_q} M_{q-1} \xrightarrow{\mu_{q-1}} \cdots \xrightarrow{\mu_1} M_0$$

with the M_i indecomposable and the μ_i irreducible in $\mathcal{D}^c(R)$, with κ_q not a retraction in $\mathcal{D}^f(R)$, and with $\mu_1 \circ \cdots \circ \mu_q \circ \kappa_q \neq 0$. I do so by induction on q .

For $q = 0$, existence of κ_0 holds by equations (4) and (5).

For $q \geq 1$, the induction gives data

$${}_R(\Sigma^i k) \xrightarrow{\kappa_{q-1}} M_{q-1} \xrightarrow{\mu_{q-1}} \cdots \xrightarrow{\mu_1} M_0.$$

Let $\tau M_{q-1} \rightarrow X_q \xrightarrow{\mu'_q} M_{q-1} \rightarrow$ be an Auslander-Reiten triangle in $\mathcal{D}^c(R)$. By [5, lem. 4.2] this is also an Auslander-Reiten triangle in $\mathcal{D}^f(R)$, so since ${}_R(\Sigma^i k) \xrightarrow{\kappa_{q-1}} M_{q-1}$ is not a retraction in $\mathcal{D}^f(R)$, it factors through X_q as ${}_R(\Sigma^i k) \xrightarrow{\kappa'_q} X_q \xrightarrow{\mu'_q} M_{q-1}$, and so $\mu_1 \circ \cdots \circ \mu_{q-1} \circ \mu'_q \circ \kappa'_q = \mu_1 \circ \cdots \circ \mu_{q-1} \circ \kappa_{q-1}$ is non-zero.

Splitting X_q into indecomposable summands and κ'_q and μ'_q into components gives that there exists ${}_R(\Sigma^i k) \xrightarrow{\kappa_q} M_q \xrightarrow{\mu_q} M_{q-1}$ with $\mu_1 \circ \cdots \circ \mu_{q-1} \circ \mu_q \circ \kappa_q \neq 0$.

Here M_q is an indecomposable summand of X_q so is in $\mathcal{D}^c(R)$, so equation (5) gives that κ_q is not a retraction in $\mathcal{D}^f(R)$. And μ_q is a component of μ'_q , so μ_q is irreducible in $\mathcal{D}^c(R)$ by [4, prop. 3.5]. $\square$

I now return to the function β .

Corollary 3.5. *Suppose that $\mathcal{D}^c(R)$ and $\mathcal{D}^c(R^{\text{op}})$ have Auslander-Reiten triangles and that ${}_R k$ is not in $\mathcal{D}^c(R)$. Let C be a component of the Auslander-Reiten quiver of $\mathcal{D}^c(R)$. Then β is unbounded on C .*

Proof. Let M_0 be an indecomposable object of $\mathcal{D}^c(R)$ for which $[M_0]$ is a vertex of C . Lemma 3.4 says that there exist arbitrarily long sequences of indecomposable objects and irreducible morphisms

$$M_q \xrightarrow{\mu_q} M_{q-1} \xrightarrow{\mu_{q-1}} \cdots \xrightarrow{\mu_1} M_0$$

in $\mathcal{D}^c(R)$ with $\mu_1 \circ \cdots \circ \mu_q \neq 0$. Each $[M_i]$ is clearly a vertex of C . So corollary 3.3 implies that β cannot be bounded on C . $\square$

4. STRUCTURE OF THE AUSLANDER-REITEN QUIVER

This section combines the material of previous sections to prove the main result of this paper, theorem 0.1.

By lemma 1.3, if $\mathcal{D}^c(R)$ and $\mathcal{D}^c(R^{\text{op}})$ have Auslander-Reiten triangles, then each component C of the Auslander-Reiten quiver of $\mathcal{D}^c(R)$

is of the form $\mathbb{Z}T/\Pi$, where T is a directed tree and $\Pi \subseteq \text{Aut}(\mathbb{Z}T)$ is an admissible group of automorphisms.

Recall from [2, sec. 4.15] that the vertices of $\mathbb{Z}T$ are the pairs (p, t) where p is an integer and t is a vertex of T , and that the vertices of $\mathbb{Z}T/\Pi$ are the orbits $\Pi(p, t)$ under the group Π . By identifying $\mathbb{Z}T/\Pi$ with C , I shall also view the $\Pi(p, t)$'s as vertices of C , so the following makes sense.

First, the start of section 2 constructs a labelling of the Auslander-Reiten quiver, and so in particular a labelling of C . Now, if there is an arrow $t \rightarrow u$ in T , then there is an arrow $\Pi(0, t) \rightarrow \Pi(0, u)$ in $\mathbb{Z}T/\Pi$, that is, an arrow $\Pi(0, t) \rightarrow \Pi(0, u)$ in C . Hence the labelling of C induces a labelling of T by

$$(a_{t \rightarrow u}, b_{t \rightarrow u}) = (a_{\Pi(0, t) \rightarrow \Pi(0, u)}, b_{\Pi(0, t) \rightarrow \Pi(0, u)}). \quad (6)$$

Secondly, the start of section 3 defines a function β on the vertices of C . That is, β is defined on the $\Pi(p, t)$'s, and so induces a function f on T by

$$f(t) = \beta(\Pi(0, t)). \quad (7)$$

Lemma 4.1. *Suppose that $D^c(R)$ and $D^c(R^{\text{op}})$ have Auslander-Reiten triangles.*

- (i) *The function f is additive with respect to the labelling of T given by equation (6).*
- (ii) *The function f is unbounded on T .*

To explain part (i), let me recall from [2, sec. 4.5] that when T is a labelled directed tree, a function f on the vertices of T is called additive if it satisfies

$$2f(t) - \sum_{s \rightarrow t} a_{s \rightarrow t} f(s) - \sum_{t \rightarrow u} b_{t \rightarrow u} f(u) = 0 \quad (8)$$

for each vertex t .

Proof of Lemma 4.1. (i) To show that the left hand side of equation (8) is zero, let me rewrite it as

$$\begin{aligned} 2\beta(\Pi(0, t)) - \sum_{s \rightarrow t} a_{\Pi(0, s) \rightarrow \Pi(0, t)} \beta(\Pi(0, s)) \\ - \sum_{t \rightarrow u} b_{\Pi(0, t) \rightarrow \Pi(0, u)} \beta(\Pi(0, u)), \end{aligned} \quad (9)$$

using equations (6) and (7).

Recall from [2, sec. 4.15] that the translation of $\mathbb{Z}T/\Pi$ is given by $\tau(\Pi(p, t)) = \Pi(p + 1, t)$. Identifying $\mathbb{Z}T/\Pi$ with C , lemma 2.1(i) gives

$b_{\Pi(0,t) \rightarrow \Pi(0,u)} = a_{\tau(\Pi(0,u)) \rightarrow \Pi(0,t)} = a_{\Pi(1,u) \rightarrow \Pi(0,t)}$, and lemma 3.1(iii) gives $\beta(\Pi(0, v)) = \beta(\tau(\Pi(0, v))) = \beta(\Pi(1, v))$. Hence (9) is

$$\begin{aligned} \beta(\tau\Pi(0, t)) + \beta(\Pi(0, t)) - \sum_{s \rightarrow t} a_{\Pi(0,s) \rightarrow \Pi(0,t)} \beta(\Pi(0, s)) \\ - \sum_{t \rightarrow u} a_{\Pi(1,u) \rightarrow \Pi(0,t)} \beta(\Pi(1, u)). \end{aligned} \quad (10)$$

To rewrite further, recall also from [2, sec. 4.15] that the arrows in $\mathbb{Z}T$ which end in $(0, t)$ are obtained as follows: There is an arrow $(0, s) \rightarrow (0, t)$ in $\mathbb{Z}T$ for each arrow $s \rightarrow t$ in T , and there is an arrow $(1, u) \rightarrow (0, t)$ in $\mathbb{Z}T$ for each arrow $t \rightarrow u$ in T .

Moreover, the canonical map $\mathbb{Z}T \rightarrow \mathbb{Z}T/\Pi$ is a so-called covering, see [2, p. 156], so the map which sends the arrow $m \rightarrow (0, t)$ to the arrow $\Pi m \rightarrow \Pi(0, t)$ is a bijection between the arrows in $\mathbb{Z}T$ which end in $(0, t)$ and the arrows in $\mathbb{Z}T/\Pi$ which end in $\Pi(0, t)$. All this implies that taken together, the two sums in (10) contain exactly one summand for each arrow in $\mathbb{Z}T/\Pi$ which ends in $\Pi(0, t)$, so (10) is

$$\beta(\tau\Pi(0, t)) + \beta(\Pi(0, t)) - \sum_{m \rightarrow \Pi(0,t)} a_{m \rightarrow \Pi(0,t)} \beta(m).$$

Identifying $\mathbb{Z}T/\Pi$ with C , this is zero by lemma 3.1(ii).

(ii) I have

$$\beta(\Pi(p, t)) = \beta(\tau^p(\Pi(0, t))) \stackrel{(a)}{=} \beta(\Pi(0, t)) = f(t),$$

where (a) is by lemma 3.1(iii). So if f were bounded on T , then β would be bounded on C which it is not by corollary 3.5. $\square$

The following is my abstract main result, which has theorem 0.1 as an easy consequence.

I emphasize that as above, C is a component of the Auslander-Reiten quiver viewed as a stable translation quiver, so C is closed under the Auslander-Reiten translation τ and its inverse, and is a connected stable translation quiver with translation the restriction of τ .

Theorem 4.2. *Suppose that $D^c(R)$ and $D^c(R^{op})$ have Auslander-Reiten triangles and that ${}_R k$ is not in $D^c(R)$. Let C be a component of the Auslander-Reiten quiver of $D^c(R)$.*

Then C is isomorphic to $\mathbb{Z}A_\infty$ as a stable translation quiver.

Proof. Lemma 1.3 says that C is of the form $\mathbb{Z}T/\Pi$.

Lemma 4.1 gives an additive function f on T , and shows that f is unbounded. Consequently, the underlying graph of T must be A_∞ by [2, thm. 4.5.8(iv)]. So C is $\mathbb{Z}A_\infty/\Pi$. (Note that when the underlying graph of T is A_∞ , the quiver $\mathbb{Z}T$ is determined up to isomorphism,

independently of the directions of the arrows in T . So it makes sense to denote $\mathbb{Z}T$ by $\mathbb{Z}A_\infty$.)

Now, Π must send vertices at the end of $\mathbb{Z}A_\infty$ to other such vertices. If Π acted non-trivially on the vertices at the end of $\mathbb{Z}A_\infty$, then there would exist a g in Π and a vertex m at the end of $\mathbb{Z}A_\infty$ so that gm was a different vertex at the end of $\mathbb{Z}A_\infty$. As the other vertices at the end of $\mathbb{Z}A_\infty$ have the form $\tau^p m$ for $p \neq 0$, this would mean $gm = \tau^p m$ for some $p \neq 0$. Then m and $\tau^p m$ would get identified in $\mathbb{Z}A_\infty/\Pi$, so Πm would be a fixed point of τ^p in $\mathbb{Z}A_\infty/\Pi$, that is, a fixed point of τ^p in C contradicting lemma 1.1(iii).

So Π acts trivially on the vertices at the end of $\mathbb{Z}A_\infty$, and it is easy to see that this forces Π to act trivially on all of $\mathbb{Z}A_\infty$. So $\mathbb{Z}A_\infty/\Pi$ is just $\mathbb{Z}A_\infty$, and the theorem follows. $\square$

Proof of Theorem 0.1. This proof uses a bit of rational homotopy theory for which my source is [3].

By [3, exam. 6, p. 146] I have that $C^*(X; k)$ is equivalent by a series of quasi-isomorphisms to a DGA, R , which satisfies setup 0.2. Hence the various derived categories of $C^*(X; k)$ are equivalent to those of R by [6, thm. III.4.2], so theorem 4.2 can be applied to $C^*(X; k)$.

To get the desired conclusion, that each component C of the Auslander-Reiten quiver of $D^c(C^*(X; k))$ is isomorphic to $\mathbb{Z}A_\infty$, I must show that the premises of theorem 4.2 hold for $C^*(X; k)$. So I must show that $D^c(C^*(X; k))$ and $D^c(C^*(X; k)^{\text{op}})$ have Auslander-Reiten triangles, and that ${}_{C^*(X; k)}k$ is not in $D^c(C^*(X; k))$. The first two claims hold by the main result of [5], theorem 6.3, because X has Poincaré duality over k . The last claim can be seen as follows.

By assumption, X has Poincaré duality of dimension $d \geq 2$ over k , so I have $H^{\geq 2}(X; k) \neq 0$. Hence the minimal Sullivan model ΛV of $C^*(X; k)$ has $V^{\geq 2} \neq 0$; see [3, chp. 12]. But V is $\text{Hom}_{\mathbb{Z}}(\pi_* X, k)$ by [3, chp. 15], where $\pi_* X$ is the sequence of homotopy groups of X , so as k has characteristic zero, $\pi_{\geq 2} X$ cannot be torsion, so $\mathbb{Q} \otimes_{\mathbb{Z}} \pi_{\geq 2} X \neq 0$. Since [3, p. 434] gives $\dim_{\mathbb{Q}}(\mathbb{Q} \otimes_{\mathbb{Z}} \pi_{\text{odd}} X) - \dim_{\mathbb{Q}}(\mathbb{Q} \otimes_{\mathbb{Z}} \pi_{\text{even}} X) \geq 0$, there follows $\mathbb{Q} \otimes_{\mathbb{Z}} \pi_{\text{odd}} X \neq 0$. This implies that the denominator is non-trivial in the formula [3, (33.7)] from which follows $\dim_{\mathbb{Q}} H^*(\Omega X; \mathbb{Q}) = \infty$, where ΩX is the Moore loop space of X . So also

$$\dim_k H^*(\Omega X; k) = \infty. \quad (11)$$

Now consider the Moore path space fibration $\Omega X \longrightarrow PX \longrightarrow X$ from [3, exam. 1, p. 29]. The Moore path space PX is contractible, and this implies that the DG left- $C^*(X; k)$ -module $C^*(PX; k)$ is quasi-isomorphic to ${}_{C^*(X; k)}k$. Inserting the fibration into the Eilenberg-Moore theorem [3, thm. 7.5] therefore gives a quasi-isomorphism $C^*(\Omega X; k) \simeq$

$k \otimes_{C^*(X;k)}^L k$, and taking cohomology shows

$$H^*(\Omega X; k) \cong \mathrm{Tor}_{-*}^{C^*(X;k)}(k, k).$$

Combining with equation (11) gives

$$\dim_k \mathrm{Tor}^{C^*(X;k)}(k, k) = \infty.$$

So ${}_{C^*(X;k)} C^*(X;k)$ cannot be finitely built from ${}_{C^*(X;k)} C^*(X;k)$ whence ${}_{C^*(X;k)} k$ is not in $D^c(C^*(X;k))$ by [5, lem. 3.2]. $\square$

REFERENCES

- [1] M. Auslander, I. Reiten, and S. Smalø, “Representation theory of Artin algebras”, Cambridge Stud. Adv. Math., Vol. 36, Cambridge University Press, Cambridge, 1997, paperback edition.
- [2] D. Benson, “Representations and cohomology, I”, Cambridge Stud. Adv. Math., Vol. 30, Cambridge University Press, Cambridge, 1998, paperback edition.
- [3] Y. Félix, S. Halperin, and J.-C. Thomas, “Rational Homotopy Theory”, Grad. Texts in Math., Vol. 205, Springer, Berlin, 2001.
- [4] D. Happel, *On the derived category of a finite-dimensional algebra*, Comment. Math. Helv. **62** (1987), 339–389.
- [5] P. Jørgensen, Auslander-Reiten triangles and quivers over topological spaces, preprint (2002). [math.AT/0304079](https://arxiv.org/abs/math/0304079).
- [6] I. Kriz and J. P. May, “Operads, algebras, modules and motives”, Astérisque **233** (1995).

DANISH NATIONAL LIBRARY OF SCIENCE AND MEDICINE, NØRRE ALLÉ 49,
2200 KØBENHAVN N, DK–DENMARK

E-mail address: pej@dnlib.dk, www.geocities.com/popjoerg